

[Published on PNAS March 13, 2007 vol. 104 no. 11](#) Mark S. Pidkowich\*, Huu Tam  
Nguyen\*,  
Heilmann  
,  
Till Ischebeck  
, and  
John Shanklin  
†

+Author Affiliations

*Department of Biology, Brookhaven National Laboratory, Upton, NY 11973*

*Edited by George H. Lorimer, University of Maryland, College Park, MD, and approved January 22, 2007 (received for review December 14, 2006)*

beta-Ketoacyl-acyl carrier protein (ACP) synthase II (KASII) elongates 16:0-ACP to 18:0-ACP in the plastid, where it competes with three other enzymes at the first major branch point in fatty acid biosynthesis. Despite its key metabolic location, the influence of KASII in determining seed oil composition remains unclear, in part because the biochemical consequences of the *fab1-1* mutation were unresolved. Thus, *fab1-1*, and a newly identified knockout allele, *fab1-2*, were analyzed in the context of the hypothesis that modulating KASII activity is sufficient to convert the composition of a temperate seed oil into that of a palm-like tropical oil. No homozygous *fab1-2* individuals were identified in progeny of self-fertilized heterozygous *fab1-2* plants, 1/4 of which aborted before the torpedo stage, suggesting that *fab1-2* represents a complete loss of function and results in lethality when homozygous. Consistent

with this hypothesis, homozygous

*fab1-2*

plants were identified when a

*fab1-1*

transgene was introduced, demonstrating that

*fab1-1*

encodes an active KASII. Strong seed-specific hairpin-RNAi reductions in

*FAB1*

expression resulted in abortion of 1/4 of the embryos in an apparent phenocopy of

*fab1-2*

homozygosity. In less severe

*FAB1*

hairpin-RNAi individuals, embryos developed normally and exhibited a 1:2:1 segregation ratio for palmitate accumulation. Thus, early embryo development appears sensitive to elevated 16:0, whereas at later stages, up to 53% of 16:0, i.e., a 7-fold increase over wild-type levels, is tolerated. These results resolve the role of KASII in seed metabolism and demonstrate that modulation of

*Arabidopsis*

KASII levels is sufficient to convert its temperate oilseed composition to that of a palm-like tropical oil.

**(\*) Mark S Pidkowich and Huu Tam Nguyen contributed equally to this work**

<http://www.pnas.org/content/104/11/4742.full>